

Colonoscopy Preparation Instructions with PEG-3350 (Trilyte, Colyte, NuLyteLy, or GoLYTELY)



IMPORTANT: Not following the instructions below may result in your procedure being cancelled and rescheduled.

1. Eating solid foods the day before your procedure may result in a suboptimal procedure.
2. **No eating or drinking anything 3 hours prior to the procedure** or your procedure will be delayed until the end of the day.
3. You need a **responsible adult age 18+** with a valid unrestricted driver's license to take you home. You are not permitted to take a taxi, bus, or service like Uber.
4. Follow **ONLY** the instructions given by your gastroenterologist or on this instruction sheet. **DO NOT follow instructions from any other source including the instructions on your prep bottle or box.**
5. If a polyp is removed we recommend no international travel for 2 weeks after your **colonoscopy** procedure. This is due to the potential risk of bleeding if large polyps are removed. Check with your gastroenterologist if you're planning to travel internationally.
6. **A successful prep is when your bowel movement is a clear yellow liquid.**

Seven (7) days before your procedure:

STOP:

1. Aspirin unless prescribed by a physician
2. NSAIDS including: Ibuprofen (Motrin, Advil), Naproxen (Aleve, Naprosyn), and similar medications
3. All seeds, nuts, popcorn, Iron supplements.

In the event you have taken any of the above, please stop taking them and let your physician know on the day of your procedure.

OKAY/APPROVED:

1. Any regular medications (especially blood pressure and/or cardiac medications)
2. Acetaminophen (Tylenol)
3. Foods such as meat (chicken, pork, beef), fish, tofu, white bread, white rice, noodles, yogurt, fruits and vegetables (remove skin and seeds), pasta, eggs, pancakes, cheese and dairy products

You will receive additional instructions, guided by your prescribing physician, if you are taking:

- ▶ Blood thinner medications such as Warfarin (Coumadin), Heparin, Lovenox, Plavix, Aggrenox, Ticlid, Pradaxa
- ▶ Diabetes medications

The ENTIRE day before your procedure:

STOP (in addition to above):

1. Any solid food
2. Dairy Products
3. Alcohol

OKAY/APPROVED:

1. Any regular medications (especially blood pressure and/or cardiac medications and acetaminophen (Tylenol))
2. Water and clear liquids including:
Juices: apple, white grape, white cranberry, lemonade
Broth: chicken, beef or vegetable
Soda: 7-Up, Coke
Other: black coffee, tea, popsicles, jello, sports drinks

*****Make sure to drink clear liquids throughout the day to keep yourself well hydrated.**

6 PM the day before your procedure: Drink half of the prep solution.

1. Add water (and flavor packet, if supplied) to the top line on the enclosed container.
2. Every 10-15 minutes drink one glass (8oz) of the solution until you have consumed half of the solution.
3. Put aside the other half of prep to take later. You can store it in the refrigerator or at room temperature.

...Continues on the back page

The day of your procedure:

- ▶ Continue your clear liquid diet.
- ▶ Make sure to drink clear liquids throughout the day to keep yourself well hydrated.
- ▶ Take your regularly prescribed morning medications (with a small sip of water) unless instructed otherwise.

5 hours before your scheduled procedure time:

- ▶ Drink the second half of the prep solution.
- ▶ Every 10 to 15 minutes drink one glass (8oz) of the solution until you have finished ***all*** the solution.

3 hours before your scheduled procedure time:

- ▶ **No eating or drinking until after your procedure or your procedure will be delayed until the end of the day.**
- ▶ A successful prep is when your bowel movement is a clear yellow liquid. **If you have any problems with your prep after hours (it doesn't work, or you throw it up, etc.) please call the gastroenterologist on call.**

* **Questions during business hours of 8am to 5:30 pm?** Please call the GI Scheduler line for your appointment center below:

San Carlos: (888)729-1890 **Palo Alto:** (888)729-1890 **Fremont:** (510)498-2148 **Mountain View:** (650)934-7554

* **Questions after business hours?** Select the option to speak to an operator at your appointment center and ask for GI Doctor on call:

San Carlos: (650)596-4130 **Palo Alto:** (650) 853-2972 **Fremont:** (510)498-2395 **Mountain View:** (650)934-7575

Facility: *(Scheduler please circle procedure location)*

Surgery Center San Carlos
301 Industrial Rd,
San Carlos, CA 94070
(First Floor)

Surgery Center of Palo Alto
795 El Camino Real
Palo Alto, CA 94301
(Jamplis Building, Level A)

Surgery Center Fremont
3120 Kearney St
Fremont, CA 94538
(First Floor)

Surgery Center Mountain View
701 E. El Camino Real
Mountain View, CA 94040
(Level 1)

Make sure to drink clear liquids throughout the day to keep yourself well hydrated.

*** PLEASE CALL YOUR PHYSICIANS OFFICE AT LEAST 7 DAYS PRIOR TO YOUR PROCEDURE IF YOU NEED TO CANCEL OR RESCHEDULE. ***

Appointment Date: _____

Arrival Time: _____ AM/PM

Doctor: _____

Procedure Time: _____ AM/PM